

# Non-suicidal self-injury prevalence, course, and association with suicidal thoughts and behaviors in two large, representative samples of US Army soldiers

## Original Article

**Cite this article:** Turner BJ, Kleiman EM, Nock MK (2018). Non-suicidal self-injury prevalence, course, and association with suicidal thoughts and behaviors in two large, representative samples of US Army soldiers. *Psychological Medicine* 1–11. <https://doi.org/10.1017/S0033291718002015>

Received: 15 December 2017

Revised: 23 June 2018

Accepted: 16 July 2018

### Key words:

Military; prevalence; risk; self-harm; soldier; suicide

### Author for correspondence:

Brianna J. Turner, E-mail: [briannat@uvic.ca](mailto:briannat@uvic.ca)

Brianna J. Turner<sup>1</sup>, Evan M. Kleiman<sup>2</sup> and Matthew K. Nock<sup>2</sup>

<sup>1</sup>Department of Psychology, University of Victoria, Victoria, BC, Canada and <sup>2</sup>Department of Psychology, Harvard University, Cambridge, MA, USA

### Abstract

**Background.** Non-suicidal self-injury (NSSI) prospectively predicts suicidal thoughts and behaviors in civilian populations. Despite high rates of suicide among US military members, little is known about the prevalence and course of NSSI, or how NSSI relates to suicidal thoughts and behaviors, in military personnel.

**Methods.** We conducted secondary analyses of two representative surveys of active-duty soldiers ( $N=21\,449$ ) and newly enlisted soldiers ( $N=38\,507$ ) from the Army Study to Assess Risk and Resilience in Servicemembers (Army STARRS).

**Results.** The lifetime prevalence of NSSI is 6.3% (1.2% 12-month prevalence) in active-duty soldiers and 7.9% (1.3% 12-month prevalence) in new soldiers. Demographic risk factors for lifetime NSSI include female sex, younger age, non-Hispanic white ethnicity, never having married, and lower educational attainment. The association of NSSI with temporally primary internalizing and externalizing disorders varies by service history (new *v.* active-duty soldiers) and gender (men *v.* women). In both active-duty and new soldiers, NSSI is associated with increased odds of subsequent onset of suicidal ideation [adjusted odds ratio (OR) = 1.66–1.81] and suicide attempts (adjusted OR = 2.02–2.43), although not with the transition from ideation to attempt (adjusted OR = 0.92–1.36). Soldiers with a history of NSSI are more likely to have made multiple suicide attempts, compared with soldiers without NSSI.

**Conclusions.** NSSI is prevalent among US Army soldiers and is associated with significantly increased odds of later suicidal thoughts and behaviors, even after NSSI has resolved. Suicide risk assessments in military populations should screen for history of NSSI.

Suicide is the 10th leading cause of death in the USA, resulting in over 40 000 deaths per year (Centers for Disease Control and Prevention, 2018). Although military personnel have historically had lower suicide rates relative to civilians, increases over the past decade have resulted in suicide rates among Servicemembers that match or exceed civilian rates (Kuehn, 2009; Armed Forces Health Surveillance Center, 2012; Nock *et al.*, 2013a, 2014). An important direction for suicide prevention is to identify US military personnel who are at high risk of attempting or dying by suicide, understand when and why they are at risk, and develop assessments and interventions that can effectively mitigate this danger (Nock, 2016).

Non-suicidal self-injury (NSSI), defined as the deliberate, self-inflicted damage of any body tissue without suicidal intent and for purposes not socially or culturally sanctioned (International Society for the Study of Self-Injury, 2018), is a strong prospective predictor of suicide attempts (SA) in civilian populations (Asarnow *et al.*, 2011; Wilkinson *et al.*, 2011; Hamza *et al.*, 2012; Ribeiro *et al.*, 2016). Yet, to our knowledge, only a handful of published studies have investigated the prevalence and characteristics of NSSI in military samples (Klonsky *et al.*, 2003; Bryan and Bryan, 2014; Kimbrel *et al.*, 2014; Shelef *et al.*, 2014; Bryan *et al.*, 2015a, 2015b; Kimbrel *et al.*, 2015, 2016, 2017; Calhoun *et al.*, 2017; Baer *et al.*, 2018). Like civilian samples, NSSI in military samples is more prevalent in women (Bryan and Bryan, 2014; cf. Bresin and Schoenleber, 2015), is typically enacted using multiple methods (Bryan and Bryan, 2014; cf. Klonsky, 2011), and is associated with elevated rates of suicidal ideation (SI) and SA (Kimbrel *et al.*, 2014; Bryan *et al.*, 2015a, 2015b; Kimbrel *et al.*, 2015; Baer *et al.*, 2018; cf. Hamza *et al.*, 2012). However, previous studies have also produced findings that diverge from civilian populations, including higher lifetime NSSI prevalence estimates (4–30% in military samples *v.* 3–6% in civilian samples; Klonsky *et al.*, 2003; Bryan *et al.*, 2015b; Kimbrel *et al.*, 2016; cf. Klonsky 2011; Plener *et al.*, 2016) and later ages of onset (17 years in military samples *v.* 12–14 years in civilians; Bryan and Bryan, 2014; cf. Whitlock and Selekman, 2014). Other important features, such as the typical course of NSSI following onset, have rarely been examined in military samples.

Previous studies are limited in several important ways. First, most studies have examined specialized samples [e.g. military personnel and veterans recruited from post-secondary

institutions (Bryan and Bryan, 2014; Bryan *et al.*, 2015a) or mental health services (Kimbrel *et al.*, 2014; Bryan *et al.*, 2015b; Baer *et al.*, 2018), or who are serving in countries with conscription (Shelef *et al.*, 2014)] that could limit the generality of the findings. Only one previous study examined NSSI in a large, representative sample of military personnel ( $N = 1986$  Air force recruits attending Basic Combat Training; Klonsky *et al.*, 2003). Moreover, about half of previous studies focus on veterans, with comparatively fewer studies examining active-duty personnel. Second, while studies support an association of NSSI with other self-injurious thoughts and behaviors (SITBs), such as SI and SA (Kimbrel *et al.*, 2015, 2016; Baer *et al.*, 2018), only a couple of studies have examined the sequencing and features of NSSI that are important for understanding trajectories among these behaviors (Bryan *et al.*, 2015a, 2015b). The current study uses two large, representative samples of US Army Soldiers to address these gaps in knowledge.

## Method

This study presents secondary analyses of two publicly available datasets from the Army Study to Assess Risk and Resilience in Servicemembers (Army STARRS; Heeringa *et al.*, 2013; Kessler *et al.*, 2013b; Ursano *et al.*, 2014): the All Army Study (AAS) and the New Soldier Study (NSS). The data sources are the paper-and-pencil or computerized self-administered questionnaires (SAQs), which were completed in group sessions to assess demographics, psychiatric symptoms, as well as history and features of SITBs. The study designs, field procedures, and sample weightings are detailed elsewhere (Heeringa *et al.*, 2013; Kessler *et al.*, 2013b, 2013c). We provide a brief summary below.

### AAS sample

The AAS uses a representative sample of the active-duty Army population across the deployment cycle and inclusive of three Army components (regular Army soldiers, activated Army National Guard, and Army Reserve soldiers). Data were collected between January 2011 and April 2013. A total of 47 744 soldiers attended study consent briefings, of whom 81% ( $n = 38\,837$ ) consented to participate and 73% ( $n = 34\,813$ ) returned completed SAQs. The final integrated sample includes 21 449 soldiers ( $n = 2504$  female,  $n = 18\,790$  male) who completed the SAQ and consented to data linkage with their administrative records.

### NSS sample

The NSS uses a representative sample of soldiers entering the Army, inclusive of regular Army, activated Army National Guard, and Army Reserve soldiers, who participated in the study during the Reception Battalion week, prior to the start of Basic Combat Training, at one of three Army installations. Data were collected between April 2011 and November 2012. A total of 58 276 new soldiers attended study consent briefings, of whom 1.2% ( $n = 698$ ) were ineligible (due to age  $<18$  years<sup>†</sup>), 95.8% ( $n = 55\,814$ ) consented to participate, and 87.1% ( $n = 50\,765$ ) returned completed SAQs. The final sample includes 38 507 soldiers ( $n = 6518$  female,  $n = 31\,445$  male) who completed the SAQ and consented to data linkage with their administrative records.

<sup>†</sup>The notes appear after the main text.

## Measures

### Demographics

Respondents self-reported their gender (male/female), age in years, race-ethnicity, nativity, marital status, educational attainment, parental educational attainment, and Army service component. AAS participants reported on their number of deployments.

### Non-suicidal self-injury

NSSI was assessed with the question ‘did you ever do something to hurt yourself on purpose, but without wanting to die (i.e. cutting yourself, hitting yourself, or burning yourself)?’ Respondents who answered in the affirmative completed follow-up questions regarding: (a) age of onset, (b) age of most recent engagement, and (c) number of lifetime NSSI episodes (with categorical response options, presented in Table 1). Ages of onset and recency were sometimes reported on a continuous scale (in years) and sometimes reported on a categorical scale for onsets under 18, depending upon the survey version. Categorically reported ages were recoded to a continuous scale (‘ $<13$  years old’ = 12, ‘13–15 years old’ = 14, and ‘16–17 years old’ = 16). Given that engagement in 10 or more acts of NSSI is associated with greater psychopathology, distress, and impairment (Muehlenkamp and Brausch, 2016), we used a cut-off of  $>10$  lifetime episodes of NSSI to designate soldiers who engaged in chronic (*v.* episodic/infrequent) NSSI.

### Mental disorders

Respondents completed an adapted version of the Composite International Diagnostic Interview Screening Scales (CIDI-SS; Kessler *et al.*, 2013a) to assess the lifetime presence and age of onset of several psychiatric disorders. The AAS assessed: major depression, bipolar disorder, panic disorder, agoraphobia, specific phobia, social phobia, generalized anxiety disorder, obsessive-compulsive disorder, intermittent explosive disorder, attention-deficit hyperactivity disorder (within the past 6 months), substance use disorders, and any other serious mental illness. The NSS assessed: major depression, bipolar disorder, panic disorder, generalized anxiety disorder, intermittent explosive disorder, attention-deficit hyperactivity disorder (within the past 6 months), oppositional defiant disorder, conduct disorder, and substance use disorders. In addition, respondents completed a screening version of the Posttraumatic Stress Disorder Checklist (PCL; Weathers *et al.*, 2013). DSM-IV diagnostic hierarchies and organic exclusion rules were not considered in assigning diagnoses. Ages of onset that were reported on a categorical scale were recoded to a continuous scale (i.e. in years) using the median value (where the median fell between whole numbers, we used the smaller value). The CIDI-SS and PCL have good concordance with diagnoses derived from clinical interviews (area under the curve = 0.69–0.79; Kessler *et al.*, 2013d).

### Suicidal thoughts and behaviors

Items assessing suicidal thoughts and behaviors were adapted from the Columbia Suicidal Severity Rating Scale (Posner *et al.*, 2011). History of SI was assessed with the question ‘did you ever in your life have thoughts of killing yourself?’<sup>2</sup>. History of SA was assessed with the question ‘did you ever make a suicide attempt (i.e. purposefully hurt yourself with at least some intention to die)?’ Respondents who endorsed SI or SA were asked follow-up questions regarding disclosure of suicidal thoughts, number of lifetime attempts, alcohol or drug use prior to attempts, and medical attention following the attempt (*medically*

**Table 1.** Lifetime and 12-month prevalence, age of onset and frequency of NSSI in the weighted AAS and NSS samples

		All Army Study (AAS)												
		Lifetime prevalence			12-month prevalence			Age of NSSI onset			Lifetime NSSI frequency			
	<i>n</i>	Weighted (%)	s.e.	<i>n</i>	Weighted (%)	s.e.	Mean	s.e.	1-2 (%)	3-5 (%)	6-10 (%)	11+ (%)		
Total sample ( <i>n</i> = 20 369)	1299	6.3	0.3	246	1.2	0.1	16.56	0.29	30.6	26.5	14.9	27.9		
Females ( <i>n</i> = 2413)	274	10.8	0.9	50	2.2	0.4	16.26	0.30	34.3	24.5	13.7	27.4		
Males ( <i>n</i> = 17 956)	1025	5.6	0.3	196	1.0	0.1	16.65	0.25	29.5	27.2	15.3	28.0		
		New Soldier Study (NSS)												
		Lifetime prevalence			12-month prevalence			NSSI onset (%)			Lifetime NSSI frequency			
	<i>n</i>	Weighted (%)	s.e.	<i>n</i>	Weighted (%)	s.e.	<13 years	13-15 years	16-17 years	>17 years	1-2 (%)	3-5 (%)	6-10 (%)	11+ (%)
Total sample ( <i>n</i> = 38 507)	2929	7.9	0.2	436	1.3	0.1	19.5	40.4	27.4	12.6	37.1	23.7	14.5	24.7
Females ( <i>n</i> = 6518)	653	10.4	0.4	62	1.1	0.1	16.5	54.9	19.9	8.7	36.1	24.1	14.3	25.5
Males ( <i>n</i> = 31 445)	2223	7.3	0.2	374	1.3	0.1	20.4	36.1	29.7	13.8	37.4	23.6	14.6	24.5

NSSI, non-suicidal self-injury.

*serious attempts* were defined as any SA that required overnight hospitalization). Consistent with the NSSI items, ages of SI or SA onset that were reported on a categorical scale were recoded to a continuous scale.

**Data analysis**


Analyses were carried out using weighted data. Design effects, weighting, and clustering were handled using the design-based Taylor series linearization method to estimate standard errors. We used cross-tabulations to estimate the prevalence, frequency, and (for categorical response options) ages of onset ranges for NSSI. We used a Cox regression to derive the cumulative hazard function (without predictors) for NSSI onset in the AAS data only, which more often used continuous response scales for NSSI onset than the NSS sample. The associations of NSSI onset with time-invariant demographic and military service variables were estimated via Cox regressions. For time-varying predictors (i.e. psychiatric disorders predicting subsequent NSSI; NSSI predicting subsequent suicidal thoughts and behaviors), we used discrete-time survival models, with person-year as the unit of analysis and a logistic link function. The person-year file was adjusted so that only ages that captured the majority (>95%) of disorder onsets were included: in the AAS, this included years 5-50; in the NSS, this included years 12-34. The optimal model for time was evaluated by including effects across the ordered ladder of polynomials until reductions in model deviance were minimized (Singer and Willett, 2003). Demographic covariates that were significantly associated with each outcome in a multivariate model were included in subsequent discrete-time survival models. To ensure that only temporally primary associations were considered, we used lagged predictor variables in the person-year file, such that changes in predictor statuses (present/absent) up to the year *prior* to the outcome were considered, but changes in the predictor that occurred during the same and subsequent years to the outcome were not included. Survival coefficients and their standard errors were exponentiated and reported as odds ratios (OR) with 95% confidence intervals (CI). Analyses were carried out in IBM SPSS Statistics version 24 using the *Complex Samples* module and SAS version 9.4 using the *proc surveylogistic* procedure.

**Results**

**Lifetime and 12-month prevalence of NSSI**

As shown in Table 1, the lifetime prevalence of NSSI is 6.3% in the AAS sample and 7.9% in the NSS sample. Stratifying by age reveals that lifetime prevalence declines with age in new soldiers (10.0% in 18-19 years old, 6.7% in 20-24 years old, 4.1% in 25+ years old), but not in active-duty soldiers (5.5% in 18-19 years old, 7.7% in 20-24 years old, and 5.8% in 25+ years old). In both samples, lifetime history of NSSI is more common in women than men (AAS: OR 2.04, 95% CI 1.64-2.53; NSS: OR 1.48, 95% CI 1.34-1.63).

The 12-month prevalence of NSSI is 1.2% (s.e. = 0.1%) in active-duty soldiers and 1.3% (s.e. = 0.1%) in new soldiers. Relative to soldiers with remote histories of NSSI, soldiers with past-year NSSI are more likely to have at least one lifetime psychiatric disorder (AAS: OR 2.16, 95% CI 1.11-4.20; NSS: OR 1.79, 95% CI 1.28-2.52) and receive mental health services in the past year (AAS: OR 2.08, 95% CI 1.18-3.67; NSS: OR 2.02, 95% CI 1.39-2.93). The average time since last NSSI episode in active-duty



**Fig. 1.** Cumulative hazard function (in bold) and fitted polynomial trend for NSSI onset in the AAS sample.

soldiers without past-year NSSI is 5.57 years (s.e. = 0.14) and is 3.81 years (s.e. = 0.07) in new soldiers. NSSI recency is stratified by age (mean years since last NSSI episode in AAS: 18–19 years old = 1.58, s.e. = 0.36; 20–24 years old = 3.45, s.e. = 0.16; 25+ years old = 5.32, s.e. = 0.20; in NSS: 18–19 years old = 2.34, s.e. = 0.08; 20–24 years old = 3.62, s.e. = 0.10; 25+ years old = 7.82, s.e. = 0.45), suggesting that average offset of NSSI occurs in early adulthood.

### *NSSI onset and course*

NSSI most often begins during adolescence, with 68% of onsets occurring before age 18 in the AAS sample and 87% of onsets occurring before age 18 in the NSS sample. Among female soldiers, NSSI onset between ages 13 and 15 is especially common, characterizing 43% and 55% of the self-injuring women in the AAS and NSS samples, respectively. The AAS data allow for a more fine-grained analysis of NSSI onset, owing to the expanded age range of the sample and use of continuous response scales. In the AAS, adult onset NSSI (after age 20) is reported by approximately 20% of respondents, whereas childhood onset (prior to age 12) is reported by 14%. [Figure 1](#) plots the hazard function for NSSI in the AAS sample, revealing stable, low rates of NSSI until age 12, a sharply increasing prevalence from age 12 through age 20, and a more gradual increase through to age 30 when the prevalence levels off.

Following NSSI onset, more than half of soldiers report fewer than five lifetime episodes of NSSI, while approximately one-quarter endorse chronic NSSI (i.e. >10 lifetime episodes; see [Table 1](#)). Soldiers with chronic NSSI are more likely to be white (AAS: OR 1.48, 95% CI 1.00–2.02; NSS: OR 1.39, 95% CI 1.04–1.87), have a high school (v. some post-secondary) education (AAS: OR 1.40, 95% CI 1.07–1.83; NSS: OR 1.43, 95% CI 1.13–1.82) and report at least one lifetime psychiatric disorder (AAS: OR 2.50, 95% CI 1.03–6.07; NSS: OR 2.42, 95% CI 1.79–3.26), but do not differ with respect to gender (AAS: OR 0.97, 95%

CI 0.66–1.43; NSS: OR 1.06, 95% CI 0.84–1.33) or marital status (AAS: OR 0.73, 95% CI 0.48–1.10; NSS: OR 1.00, 95% CI 0.73–1.35) compared with soldiers with ten or fewer NSSI episodes. New soldiers with chronic NSSI episodes are younger than those with ten or fewer lifetime NSSI episodes (NSS: OR 0.92, 95% CI 0.89–0.96), but this is not true of active-duty soldiers (AAS: OR 0.99, 95% CI 0.97–1.01). Whereas active-duty soldiers with chronic NSSI are more likely to have received mental health services in the past 12 months (AAS: OR 2.02, 95% CI 1.34–3.04), this was not true of new soldiers (NSS: OR 1.02, 95% CI 0.76–1.38).

### *Demographic risk factors for NSSI*

The odds of having a lifetime history of NSSI are elevated among women, soldiers who were <25 years old, and soldiers who had never been married ([Table 2](#)). Soldiers who self-identify as non-Hispanic black or Hispanic have lower odds of NSSI compared with soldiers who self-identify as non-Hispanic white. Among active-duty soldiers, those who have been deployed three or more times and those whose parents had not completed high school have lower odds of having engaged in NSSI. Among new soldiers, those who immigrated to the USA have lower odds of NSSI, whereas those with less educational attainment (<4 years post-secondary) have higher odds of NSSI. Whereas active-duty soldiers who are serving in the Army Reserves National Guard or Army Reserves have lower odds of engaging in NSSI, relative to regular Army soldiers, new soldiers who enlist in the Army Reserves National Guard or General Reserves have higher odds of NSSI than those who enlist in the regular Army.

### *Psychiatric risk factors for NSSI*

A number of psychiatric disorders are associated with elevated odds of subsequent NSSI onset ([Table 3](#)). Among active-duty

**Table 2.** Results of the weighted Cox proportional-hazards regressions assessing the associations of sociodemographic and military service characteristics with lifetime and 12-month NSSI

	All Army Study (AAS)				New Soldier Study (NSS)			
	Lifetime NSSI		12-month NSSI (among lifetime)		Lifetime NSSI		12-month NSSI (among lifetime)	
	OR	95% CI	OR	95% CI	OR	95% CI	OR	95% CI
<b>Sex</b>								
Men	1 (Reference)		1 (Reference)		1 (Reference)		1 (Reference)	
Women	<b>1.99</b>	<b>1.61–2.47</b>	<b>1.51</b>	<b>1.10–2.07</b>	<b>1.47</b>	<b>1.34–1.62</b>	<b>0.59</b>	<b>0.44–0.78</b>
<b>Age</b>								
≥25	1 (Reference)		1 (Reference)		1 (Reference)		1 (Reference)	
20–24	<b>1.64</b>	<b>1.31–2.05</b>	N/A	N/A	<b>1.95</b>	<b>1.60–2.38</b>	N/A	N/A
18–19	1.35	0.84 to 2.18	N/A	N/A	<b>3.31</b>	<b>2.74–4.00</b>	N/A	N/A
<b>Race/ethnicity</b>								
Non-Hispanic white	1 (Reference)		1 (Reference)		1 (Reference)		1 (Reference)	
Non-Hispanic black	<b>0.35</b>	<b>0.23–0.52</b>	1.36	0.83–2.22	<b>0.44</b>	<b>0.38–0.51</b>	0.74	0.51–1.06
Hispanic	<b>0.75</b>	<b>0.62–0.91</b>	0.78	0.44–1.37	<b>0.75</b>	<b>0.66–0.84</b>	0.91	0.65–1.27
All others	0.91	0.64–1.28	0.56	0.29–1.10	0.98	0.82 to 1.17	0.99	0.70–1.41
<b>Nativity</b>								
Second+ generation	1 (Reference)		1 (Reference)		1 (Reference)		1 (Reference)	
Immigrant	0.99	0.70–1.39	<b>0.28</b>	<b>0.14–0.57</b>	<b>0.69</b>	<b>0.58–0.83</b>	1.11	0.74–1.65
First generation	0.93	0.76–1.14	1.06	0.70–1.61	0.92	0.82–1.02	0.72	0.47–1.11
<b>Marital status</b>								
Married	1 (Reference)		1 (Reference)		1 (Reference)		1 (Reference)	
Never married	<b>1.27</b>	<b>1.00–1.61</b>	<b>2.49</b>	<b>1.69–3.67</b>	<b>1.43</b>	<b>1.25–1.63</b>	<b>2.07</b>	<b>1.37–3.12</b>
Previously married	1.08	0.80–1.45	1.37	0.87–2.17	1.25	0.90–1.74	1.16	0.50–2.71
<b>Education</b>								
Bachelors or higher	1 (Reference)		1 (Reference)		1 (Reference)		1 (Reference)	
Completed high school	1.23	0.99–1.52	8.38	4.82–14.21	<b>2.22</b>	<b>1.77–2.78</b>	<b>5.09</b>	<b>2.95–8.80</b>
At least some college	1.13	0.86–1.48	6.48	3.91–10.73	<b>1.80</b>	<b>1.47–2.20</b>	<b>2.01</b>	<b>1.11–3.66</b>
<b>Parental education</b>								
Bachelors or higher	1 (Reference)		1 (Reference)		1 (Reference)		1 (Reference)	
Less than high school	<b>0.58</b>	<b>0.39–0.86</b>	<b>1.95</b>	<b>1.02–3.72</b>	1.16	0.97–1.28	0.90	0.60–1.34
Completed high school	0.81	0.65–1.00	1.31	0.74–2.33	0.97	0.88–1.08	1.29	0.98–1.69
At least some college	0.95	0.74–1.23	1.35	0.78–2.34	1.06	0.95–1.19	1.10	0.81–1.49
<b>Deployments</b>								
0	1 (Reference)		1 (Reference)		1 (Reference)		1 (Reference)	
1	0.81	0.49–1.33	0.79	0.34–1.87	N/A	N/A	N/A	N/A
2	0.95	0.67–1.36	0.74	0.46–1.18	N/A	N/A	N/A	N/A
≥3	<b>0.65</b>	<b>0.46–0.93</b>	<b>0.31</b>	<b>0.20–0.47</b>	N/A	N/A	N/A	N/A
<b>Army sector</b>								
Regular Army	(Reference)		(Reference)		(Reference)		(Reference)	
ARNG	<b>0.51</b>	<b>0.33–0.78</b>	1.18	0.68–2.03	<b>1.15</b>	<b>1.02–1.29</b>	0.93	0.64–1.35

ARNG, Army Reserves/Army National Guard.

Bold text indicates statistically significant effect at  $p < 0.05$ .

**Table 3.** Results of the multivariate weighted discrete-time survival analyses assessing the associations of temporally primary psychiatric disorders with first onset of NSSI in the total samples, in men and in women

	All Army Study (AAS) <sup>a</sup>					
	Total sample		Men		Women <sup>b</sup>	
	OR	95% CI	OR	95% CI	OR	95% CI
MDE	<b>1.68</b>	<b>1.09–2.58</b>	1.63	0.92–2.88	<b>1.70</b>	<b>1.18–2.46</b>
Bipolar	1.56	0.94–2.59	1.54	0.79–3.03	1.65	0.91–2.99
Panic	1.03	0.64–1.67	0.86	0.51–1.45	1.70	0.56–5.19
Agoraphobia	0.83	0.45–1.52	0.89	0.44–1.81	0.73	0.33–1.64
Specific phobia	1.09	0.80–1.48	1.05	0.71–1.54	1.22	0.84–1.78
Social phobia	<b>1.80</b>	<b>1.45–2.24</b>	<b>1.92</b>	<b>1.47–2.51</b>	1.42	0.94–2.15
GAD	1.15	0.81–1.64	1.26	0.83–1.92	0.87	0.53–1.45
PTSD	0.93	0.68–1.29	0.87	0.57–1.33	1.11	0.68–1.83
OCD	<b>1.89</b>	<b>1.44–2.49</b>	<b>1.71</b>	<b>1.22–2.38</b>	<b>2.36</b>	<b>1.49–3.73</b>
IED	<b>2.35</b>	<b>1.94–2.84</b>	<b>2.22</b>	<b>1.79–2.75</b>	<b>2.73</b>	<b>1.76–4.24</b>
ADHD	<b>2.17</b>	<b>1.56–3.02</b>	<b>2.23</b>	<b>1.57–3.16</b>	<b>2.32</b>	<b>1.22–4.42</b>
SUD	1.15	0.89–1.48	1.15	0.89–1.50	1.04	0.47–2.31
SMI	1.33	0.94–1.88	<b>1.56</b>	<b>1.06–2.30</b>	0.83	0.37–1.86
	New Soldier Study (NSS) <sup>c</sup>					
	Total sample		Men		Women	
	OR	95% CI	OR	95% CI	OR	95% CI
MDE	1.02	0.84–1.25	1.05	0.82–1.35	0.98	0.63–1.51
Bipolar	0.91	0.68–1.22	1.00	0.73–1.37	0.67	0.33–1.35
Panic	1.25	0.99–1.57	1.25	0.94–1.67	1.24	0.77–2.00
GAD	0.87	0.69–1.11	0.92	0.70–1.19	0.81	0.51–1.29
PTSD	<b>2.07</b>	<b>1.84–2.34</b>	<b>1.97</b>	<b>1.73–2.24</b>	<b>2.29</b>	<b>1.77–2.96</b>
IED	<b>1.24</b>	<b>1.08–1.42</b>	<b>1.30</b>	<b>1.13–1.49</b>	1.00	0.68–1.46
ADHD	<b>1.82</b>	<b>1.59–2.08</b>	<b>1.86</b>	<b>1.58–2.19</b>	<b>1.67</b>	<b>1.25–2.25</b>
ODD	<b>2.43</b>	<b>2.14–2.76</b>	<b>2.37</b>	<b>2.05–2.73</b>	<b>2.73</b>	<b>2.10–3.55</b>
CD	<b>1.53</b>	<b>1.27–1.83</b>	<b>1.55</b>	<b>1.28–1.88</b>	1.43	0.89–2.28
SUD	<b>1.85</b>	<b>1.65–2.08</b>	<b>1.87</b>	<b>1.65–2.12</b>	<b>1.82</b>	<b>1.41–2.24</b>

MDE, major depressive episode; GAD, generalized anxiety disorder; PTSD, post-traumatic stress disorder; OCD, obsessive-compulsive disorder; IED, intermittent explosive disorder; ADHD, attention-deficit hyperactivity disorder; ODD, oppositional defiant disorder; CD, conduct disorder; SUD, substance use disorder; SMI, serious mental illness.

Bold text indicates statistically significant effect at  $p < 0.05$  in multivariate model.

<sup>a</sup>The AAS discrete-time survival model covaries for time (modeled via polynomial terms to the fourth power), sex, race, and service component (ARNG v. regular Army).

<sup>b</sup>The AAS discrete-time survival model for women removes the effect of the fourth power for time to improve model fit.

<sup>c</sup>The NSS discrete-time survival model covaries for time (modeled via polynomial terms to the third power), sex, age, and race.

soldiers, major depressive, social phobia, obsessive-compulsive, intermittent explosive, and attention-deficit hyperactivity disorders are each uniquely associated with subsequent NSSI onset in a multivariate model. Sensitivity analyses show that social phobia and serious mental illness are significantly associated with subsequent NSSI only in male active-duty soldiers, whereas major depression is significantly associated with subsequent NSSI only in female active-duty soldiers. Among new soldiers, post-traumatic stress, intermittent explosive, attention-deficit hyperactivity, oppositional defiant, conduct, and substance use disorders are each uniquely associated with subsequent NSSI in multivariate models. Sensitivity analyses show that intermittent explosive and conduct

disorder are significantly associated with subsequent NSSI only in new male soldiers.

#### *NSSI as a risk factor for suicidal behavior*

Nearly 50% of soldiers with a history of NSSI also report SI, and approximately one in six have attempted suicide, contrasting the much lower rates of ideation (8.7%) and attempt (1.8%) in soldiers without NSSI (Table 4). Lifetime NSSI is associated with significantly higher odds of a lifetime history of SI, SA, attempts among individuals with suicidal thoughts, and multiple attempts in bivariate cross-tabulations (Table 4). Among

**Table 4.** Prevalence and bivariate associations of lifetime suicidal thoughts and behaviors in soldiers with and without lifetime history of NSSI

	All Army Study (AAS)					
	Prevalence in soldiers without NSSI		Prevalence in soldiers with NSSI			
	%	S.E.	%	S.E.	OR	95% CI
Suicidal ideation	<b>8.7</b>	<b>0.3</b>	<b>48.9</b>	<b>2.4</b>	<b>10.11</b>	<b>8.38–12.20</b>
Suicide attempt	<b>1.8</b>	<b>0.1</b>	<b>17.1</b>	<b>1.4</b>	<b>11.30</b>	<b>9.02–14.15</b>
Among soldiers with SI						
Attempts	<b>18.7</b>	<b>1.4</b>	<b>33.5</b>	<b>2.5</b>	<b>2.19</b>	<b>1.69–2.83</b>
Disclosed suicidal thoughts	<b>3.9</b>	<b>0.3</b>	<b>23.1</b>	<b>2.0</b>	<b>7.42</b>	<b>5.86–9.39</b>
Among soldiers with SA						
Multiple attempts	<b>30.9</b>	<b>3.0</b>	<b>50.5</b>	<b>5.4</b>	<b>2.28</b>	<b>1.38–3.76</b>
Alcohol/drugs prior to attempt	<b>16.1</b>	<b>2.0</b>	<b>36.7</b>	<b>4.2</b>	<b>3.03</b>	<b>1.90–4.83</b>
Medically serious attempt	6.1	1.3	5.7	1.7	0.93	0.45–1.93
New Soldier Study (NSS)						
	%	S.E.	%	S.E.	OR	95% CI
Suicidal ideation	<b>6.3</b>	<b>0.2</b>	<b>49.5</b>	<b>1.2</b>	<b>14.50</b>	<b>13.13–16.02</b>
Suicide attempt	<b>0.8</b>	<b>0.1</b>	<b>14.1</b>	<b>0.7</b>	<b>20.91</b>	<b>17.93–24.38</b>
Among soldiers with SI						
Attempts	<b>10.0</b>	<b>0.7</b>	<b>25.3</b>	<b>1.2</b>	<b>3.05</b>	<b>2.55–3.65</b>
Disclosed suicidal thoughts	46.8	2.1	51.9	2.2	1.23	0.97–1.55
Among soldiers with SA						
Multiple attempts	<b>42.9</b>	<b>3.3</b>	<b>65.9</b>	<b>2.8</b>	<b>2.57</b>	<b>1.81–3.65</b>
Alcohol/drugs prior to attempt	24.4	2.8	27.3	2.6	1.16	0.75–1.80
Medically serious attempt	5.4	1.5	7.6	1.6	1.42	0.67–3.03

NSSI, non-suicidal self-injury; SI, suicidal ideation; SA, suicide attempt(s). Bold text indicates statistically significant effect at  $p < 0.05$ .

active-duty soldiers, a lifetime history of NSSI is also associated with greater odds of having used alcohol or drugs before a SA and having disclosed suicidal thoughts. Chronic NSSI (>10 episodes) is associated with elevated odds of SA (AAS: OR 2.75, 95% CI 1.86–4.06; NSS: OR 2.94, 95% CI 2.29–3.77), of making multiple attempts in new soldiers (NSS: OR 1.60, 95% CI 1.01–2.53), and of disclosing suicidal thoughts in active-duty soldiers (AAS: OR 1.61, 95% CI 1.02–2.56), relative to soldiers with infrequent/episodic NSSI. Earlier NSSI onset (before age 16) is not associated with odds of lifetime SI (AAS: OR 0.91, 95% CI 0.67–1.24; NSS: OR 0.98, 95% CI 0.83–1.17) nor with lifetime SA in active-duty soldiers (OR 0.92, 95% CI 0.65–1.29), but is associated with higher odds of lifetime attempt in new soldiers (OR 1.46, 95% CI 1.10–1.93).

In terms of sequencing, retrospective reports show that NSSI precedes ideation in 15.9–26.7% of soldiers, and precedes attempts in 22.6–41.4% of soldiers (Table 5). More commonly, NSSI and suicidal thoughts or behaviors have an onset within the same year (47.0–58.4% of soldiers with both NSSI and ideation, 34.7–58.2% of soldiers with both NSSI and attempts). Discrete-time survival models, which considered only NSSI that occurred at least 1 year prior to the onset of SI or SA, show that NSSI is associated with elevated odds of subsequent SI (bivariate OR = 3.24–4.71, adjusted OR = 1.66–1.81) and SA

(bivariate OR = 5.51–7.57, adjusted OR = 2.02–2.43), but not with attempts among soldiers with suicidal thoughts (bivariate OR = 0.88–1.73, adjusted OR = 0.92–1.10).

## Discussion

To our knowledge, this study is the largest and most representative investigation of NSSI in US military personnel to date. There are four major findings from this study. First, approximately 6–8% of US Army soldiers have a lifetime history of NSSI, while about 1% have engaged in NSSI in the past year. Second, most NSSI begins in adolescence, occurs infrequently (i.e.  $\leq 10$  lifetime episodes), and has resolved before soldiers enter military service. Nonetheless, any NSSI history is associated with elevated risk of suicidal thoughts and behaviors, and therefore merits clinical attention. Third, NSSI is over-represented in certain groups of soldiers, including those who identify as women, under 25 years old, non-Hispanic white, and never having married, as well as those with histories of internalizing or externalizing psychiatric disorders. Fourth, rates of SA are approximately 10 times higher in soldiers with a history of NSSI compared with those without this behavior. Moreover, NSSI significantly predicts subsequent onset of suicidal thoughts and behaviors, even after controlling for demographic and

**Table 5.** Results of the weighted discrete-time survival analyses assessing the associations of temporally primary NSSI and subsequent suicidal behaviors

	All Army Study (AAS)							
	Timing of NSSI relative to suicidal behavior				Discrete-time survival models			
	NSSI first		NSSI concurrent		Unadjusted		Adjusted	
	%	s.e.	%	s.e.	OR	95% CI	OR	95% CI
Suicidal ideation	26.7	2.8	47.0	2.8	<b>4.71</b>	<b>3.94–5.64</b>	<b>1.97</b>	<b>1.47–2.68<sup>a</sup></b>
Suicide attempt	41.4	4.2	34.7	3.9	<b>7.57</b>	<b>5.21–11.01</b>	<b>2.02</b>	<b>1.17–3.48<sup>b</sup></b>
Attempts in soldiers with SI	41.6	4.5	35.1	4.2	<b>1.73</b>	<b>1.18–2.52</b>	1.10	0.71–1.71 <sup>c</sup>
	New Soldier Study (NSS)							
	NSSI first		NSSI concurrent		Unadjusted		Adjusted	
	%	s.e.	%	s.e.	OR	95% CI	OR	95% CI
Suicidal ideation	15.9	1.3	58.4	1.8	<b>3.24</b>	<b>2.74–3.82</b>	<b>1.66</b>	<b>1.21–2.28<sup>d</sup></b>
Suicide attempt	22.6	3.1	58.2	4.0	<b>5.51</b>	<b>4.05–7.51</b>	<b>2.43</b>	<b>1.67–3.55<sup>e</sup></b>
Attempts in soldiers with SI	23.2	3.2	58.8	4.0	0.88	0.64–1.19	0.92	0.64–1.27 <sup>f</sup>

NSSI, non-suicidal self-injury; SI, suicidal ideation.

<sup>a</sup>The AAS multivariate survival model for suicidal ideation covaries for time (modeled via polynomial terms to the fifth power), sex, race, service component (Army Reserve/National Guard v. regular Army), and all psychiatric disorders.

<sup>b</sup>The AAS multivariate survival model for suicide attempts in the total sample covaries for time (modeled via polynomial terms to the fourth power), sex, service component (Army Reserve/National Guard v. regular Army), and all psychiatric disorders.

<sup>c</sup>The AAS multivariate survival model for suicide attempts among soldiers with SI covaries for time (modelled via polynomial terms to the fourth power), sex, service component (Army Reserve/National Guard v. regular Army), and all psychiatric disorders.

<sup>d</sup>The NSS survival model for suicidal ideation covaries for time (modeled via polynomial terms to the fourth power), sex, age, enlisted service component (Army Reserve/National Guard v. regular Army), and all psychiatric disorders.

<sup>e</sup>The NSS survival model for suicidal ideation in the total sample covaries for time (modeled via polynomial terms to the third power), sex, age, educational attainment, and all psychiatric disorders.

<sup>f</sup>The NSS survival model for suicidal attempts in the total sample covaries for time (modeled via polynomial terms to the third power), sex, age, educational attainment, and all psychiatric disorders.

psychiatric risk factors. These findings underscore the importance of including NSSI in suicide risk assessments.

Results from this study align with what we know about NSSI in civilian populations in several important ways. Our results show a similar lifetime prevalence of NSSI (6–8%) as representative studies of civilian adults (3–6%; Klonsky, 2011; Plener *et al.*, 2016), suggesting that NSSI is at least as common in Servicemembers as in civilians, and deserving of expanded research and clinical attention. Also consistent with civilian studies, the estimated prevalence of NSSI is considerably lower in the general population than in samples recruited from post-secondary institutions (12–14%; Bryan and Bryan, 2014; Bryan *et al.*, 2015a) and clinical settings (30–56%; Kimbrel *et al.*, 2014; Bryan *et al.*, 2015b; cf. Whitlock *et al.*, 2011; Muehlenkamp *et al.*, 2012a). With respect to the onset, course, and topography of NSSI, our results converge with civilian studies showing that adolescence is a high-risk period for the onset of NSSI (Barrocas *et al.*, 2012), that NSSI often resolves in late adolescence and early adulthood (Whitlock and Selekman, 2014), and that most people who engaged in NSSI do so only a few times, while a clinically important minority of individuals engage in chronic, repetitive NSSI (Muehlenkamp and Gutierrez, 2007; Zetterqvist *et al.*, 2013; Muehlenkamp and Brausch, 2016). With respect to demographic factors, our results align with civilian studies showing that women are more likely to engage in NSSI compared with men (Bresin and Schoenleber, 2015), that NSSI may be more prevalent in younger v. older cohorts (Klonsky, 2011), and that NSSI is more common in unmarried adults compared with their married peers (Klonsky, 2011). Moreover, our results also show that NSSI is associated with a range of internalizing and externalizing psychopathology,

consistent with civilian samples (Nock *et al.*, 2006, 2010). Finally, our results underscore strong associations between NSSI and range of SITBs, including SI, SA, and multiple attempts, particularly among individuals more than 10 lifetime episodes of NSSI, as in civilian samples (Whitlock *et al.*, 2008). It is worth noting that our models adopted a conservative approach by including only NSSI with an onset at least 1 year prior to suicidal behaviors, which restricted analyses to the 23–41% of SA. The lack of clear developmental sequencing of NSSI preceding SI and attempts is inconsistent with a previous military study (Bryan *et al.*, 2015b), and may be due in part to the forced-choice response options in several subsamples. Despite this limitation, our results suggest that NSSI is an important predictor of onset of later SI and attempts, but not of the ideation-to-attempt transition, in military personnel. Reconciling this finding with current theories of suicide risk (e.g. Joiner *et al.*, 2012) is an important direction for future work (see also Nock *et al.*, 2018).

In addition to highlighting similarities in NSSI presentation in military and civilian populations, our results may have practical implications for the Army. For instance, our results show that, for most Servicemembers, NSSI has discontinued by the time of enlistment, and will present as an active area of clinical concern in a minority (~1%) of enlisted and active-duty soldiers. Nonetheless, soldiers with histories of chronic NSSI, defined here as more than 10 lifetime episodes, are at elevated risk for suicidal behaviors. Assessing history and frequency of NSSI behavior in routine mental health screenings, including in pre-enlistment and pre-deployment screenings, could identify soldiers who may benefit from additional support. However, policies restricting enlistment or service duties of individuals with mental health problems (e.g. the

inclusion of 'self-mutilation' as a non-waiverable condition) may result in reticence to disclose NSSI in military settings. Given that NSSI has typically resolved by enlistment, permitting accession waivers for self-mutilation may increase disclosures and help to identify at-risk soldiers at the outset of their military careers. Previous studies suggest that waived pre-enlistment medical conditions were not associated with shorter deployment durations (Gubata *et al.*, 2013). Expanding predeployment screenings to include NSSI may be helpful in linking deployed soldiers to ongoing support while in-theatre (Warner *et al.*, 2011). Moreover, screening for NSSI in non-deployed personnel could help mitigate the high risk of suicide among soldiers who are never deployed (Schoenbaum *et al.*, 2014). Given the personal demands associated with military service, understanding the impact of NSSI on Servicemembers' occupational and mental health functioning, as well as trainings or policies that could bolster resilience, is an important direction for future work.

Findings should be interpreted in light of several limitations. First, these data are cross-sectional and retrospective in nature, and are subject to recall biases. Prospective studies that focus on adolescence, when SITBs typically begin (Nock *et al.*, 2009, 2013b), and retrospective studies using structured interviews to reduce retrospective biases (e.g. the timeline follow-back; Sobell and Sobell, 1992; see also Bagge *et al.*, 2013), can help to the temporal sequencing of SITB and the role of NSSI in subsequent SITB. Second, given the sensitive and potentially stigmatizing nature of SITBs, it is possible that soldiers under-reported their engagement in NSSI. The clarifying examples of NSSI provided in these surveys ('cutting yourself, hitting yourself, or burning yourself') may not represent the experiences of NSSI in men (Green and Jakupcak, 2016) or military personnel (Kimbrel *et al.*, 2014), potentially resulting in further under-reporting. Third, while the CIDI-SS provides an efficient and accurate method for assessing a range of psychopathology (Kessler *et al.*, 2013d), limitations remain in using self-report methods to assess psychiatric diagnoses as items cannot be clarified and responses cannot be probed. Moreover, several diagnoses that were not assessed in the AAS and NSS studies (e.g. personality disorders, eating disorders) have established relationships to NSSI (Turner *et al.*, 2015; Muehlenkamp *et al.*, 2012b), and should be included in future epidemiological studies of NSSI. The present estimates of the association between psychopathology and NSSI may be considered conservative as a result.

These limitations notwithstanding this study is the first to use multiple representative samples to investigate the prevalence, course, and consequences of NSSI in US military personnel. For the most part, results from the two samples converge, although a number of divergent findings require clarification in future studies (e.g. age and cohort effects in new *v.* active-duty soldiers; prevalence of NSSI in different service components of the Army; associations of NSSI with specific psychiatric disorders [post-traumatic stress disorder, substance use] and contextual SITB factors [alcohol/drug use]). Nonetheless, our results should increase attention to this clinically important behavior in military populations, and underscore the importance of including NSSI in comprehensive suicide risk assessments.

## Notes

<sup>1</sup> Individuals are eligible to enlist in the US Army at age 17 with parental consent, or ages 18+ without parental consent. However, approved study procedures required participants to be at least 18 years old.

<sup>2</sup> Note that for the purposes of this study, we did not include passive ideation ('did you ever wish you were dead or would go to sleep and never wake up?') in our definition of suicidal ideation. Supplementary analyses assessing the relationship of NSSI to passive ideation are available from the first author.

**Acknowledgements.** This publication is based on public use data from the Army Study to Assess Risk and Resilience in Servicemembers (Army STARRS). The data are available from the Inter-university Consortium for Political and Social Research (ICPSR) at the University of Michigan (<http://doi.org/10.3886/ICPSR35197-v1>). Army STARRS was funded by the US National Institute of Mental Health (grant number U01MH087981). The contents of this publication are solely the responsibility of the authors and do not necessarily represent the views of the Army STARRS investigators, funders, Department of the Army, or Department of Defense.

## References

- Armed Forces Health Surveillance Center (AFHSC)** (2012) Deaths while on active-duty in the U.S. Armed Forces, 1990–2011. *MSMR: Medical Surveillance Monthly Report* **19**, 2–5.
- Asarnow JR, Porta G, Spirito A, Emslie G, Clarke G, Wagner KD, Vitiello B, Keller M, Birmaher B, McCracken J, Mayes T, Berk M and Brent DA** (2011) Suicide attempts and nonsuicidal self-injury in the Treatment of Resistant Depression in Adolescents: findings from the TORDIA study. *Journal of the American Academy of Child & Adolescent Psychiatry* **50**, 772–781.
- Baer MM, LaCroix JM, Browne JC, Hassen HO, Perera KU, Weaver J, Soumoff A and Ghahramanlou-Holloway M** (2018) Non-suicidal self-injury elevates suicide risk among United States military personnel with lifetime attempted suicide. *Archives of Suicide Research* **22**, 453–464.
- Bagge CL, Glenn CR and Lee HJ** (2013) Quantifying the impact of recent negative life events on suicide attempts. *Journal of Abnormal Psychology* **122**, 359–368.
- Barrocas AL, Hankin BL, Young JF and Abela JRZ** (2012) Rates of nonsuicidal self-injury in youth: age, sex, and behavioral methods in a community sample. *Pediatrics* **130**, 39–45.
- Bresin K and Schoenleber M** (2015) Gender differences in the prevalence of nonsuicidal self-injury: a meta-analysis. *Clinical Psychology Review* **38**, 55–64.
- Bryan CJ and Bryan AO** (2014) Nonsuicidal self-injury among a sample of United States military personnel and veterans enrolled in college classes. *Journal of Clinical Psychology* **70**, 874–885.
- Bryan CJ, Rudd MD, Wertemberger E, Young-McCaughon S and Peterson A** (2015a) Nonsuicidal self-injury as a prospective predictor of suicide attempts in a clinical sample of military personnel. *Comprehensive Psychiatry* **59**, 1–7.
- Bryan CJ, Bryan AO, May AM and Klonsky ED** (2015b) Trajectories of suicide ideation, nonsuicidal self-injury, and suicide attempts in a nonclinical sample of military personnel and veterans. *Suicide & Life-threatening Behavior* **45**, 315–325.
- Calhoun PS, Van Voorhees EE, Elbogen EB, Dedert EA, Clancy CP, Hair LP, Hertzberg M, Beckham JC and Kimbrel NA** (2017) Nonsuicidal self-injury and interpersonal violence in U.S. veterans seeking help for posttraumatic stress disorder. *Psychiatry Research* **247**, 250–256.
- Centers for Disease Control and Prevention** (2018) Vital signs: trends in state suicide rates – United States, 1999–2016 and circumstances contributing to suicide – 27 states, 2015. *Morbidity and Mortality Weekly Report* **67**, 617–624.
- Green JD and Jakupcak M** (2016) Masculinity and men's self-harm behaviors: implications for non-suicidal self-injury disorder. *Psychology of Men and Masculinity* **17**, 147–155.
- Gubata ME, Oetting AA, Niebuhr DW and Cowan DN** (2013) Accession medical waivers and deployment duration in the U.S. Army. *Military Medicine* **178**, 625–630.

- Hamza CA, Stewart SL and Willoughby T (2012) Examining the link between nonsuicidal self-injury and suicidal behavior: a review of the literature and an integrated model. *Clinical Psychology Review* 32, 482–495.
- Heeringa SG, Gebler N, Colpe LJ, Fullerton CS, Hwang I, Kessler RC, Naifeh JA, Nock MK, Sampson NA, Schoenbaum M, Zaslavsky AM, Stein MB and Ursano RJ (2013) Field procedures in the Army Study to Assess Risk and Resilience in Servicemembers (Army STARRS). *International Journal of Methods in Psychiatric Research* 22, 276–287.
- International Society for the Study of Self-injury (2018) About self-injury: What is nonsuicidal self-injury? Available at <https://itriples.org/category/about-self-injury/> (Accessed 22 June 2018).
- Joiner TE, Ribeiro JD and Silva C (2012) Nonsuicidal self-injury, suicidal behavior, and their co-occurrence as viewed through the lens of the interpersonal theory of suicide. *Current Directions in Psychological Science* 21, 342–347.
- Kessler RC, Calabrese JR, Farley PA, Gruber MJ, Jewell MA, Katon W, Keck PE, Nierenberg AA, Sampson NA, Shear MK, Shillington AC, Stein MB, Thase ME and Wittchen HU (2013a) Composite International Diagnostic Interview screening scales for DSM-IV anxiety and mood disorders. *Psychological Medicine* 43, 1625–1637.
- Kessler RC, Colpe LJ, Fullerton CS, Gebler N, Naifeh JA, Nock MK, Sampson NA, Schoenbaum M, Zaslavsky AM, Stein MB, Ursano RJ and Heeringa SG (2013b) Design of the Army Study to Assess Risk and Resilience in Servicemembers (Army STARRS). *International Journal of Methods in Psychiatric Research* 22, 267–275.
- Kessler RC, Heeringa SG, Colpe LJ, Fullerton CS, Gebler N, Hwang I, Naifeh JA, Nock MK, Sampson NA, Schoenbaum M, Zaslavsky AM, Stein MB and Ursano RJ (2013c) Response bias, weighting adjustments, and design effects in the Army Study to Assess Risk and Resilience in Servicemembers (Army STARRS). *International Journal of Psychiatric Research* 22, 288–302.
- Kessler RC, Santiago PN, Colpe LJ, Dempsey CL, First MB, Heeringa SG, Stein MB, Fullerton CS, Gruber MJ, Naifeh JA, Nock MK, Sampson NA, Schoenbaum M, Zaslavsky AM and Ursano RJ (2013d) Clinical reappraisal of the Composite International Diagnostic Interview Screening Scales (CIDI-SC) in the Army Study to Assess Risk and Resilience in Servicemembers (Army STARRS). *International Journal of Methods in Psychiatric Research* 22, 303–321.
- Kimbrel NA, Johnson ME, Clancy C, Hertzberg M, Collie C, Van Voorhees EE, Dennis MF, Calhoun PS and Beckham JC (2014) Deliberate self-harm and suicidal ideation among male Iraq/Afghanistan-era veterans seeking treatment for PTSD. *Journal of Traumatic Stress* 27, 474–477.
- Kimbrel NA, Gratz KL, Tull MT, Morissette SB, Meyer EC, DeBeer BB, Silvia PJ, Calhoun PC and Beckham JC (2015) Non-suicidal self-injury as a predictor of active and passive suicidal ideation among Iraq/Afghanistan war veterans. *Psychiatry Research* 227, 360–362.
- Kimbrel NA, DeBeer BB, Meyer EC, Gulliver SB and Morissette SB (2016) Nonsuicidal self-injury and suicide attempts in Iraq/Afghanistan war veterans. *Psychiatry Research* 243, 232–237.
- Kimbrel NA, Wilson LC, Mitchell JT, Meyer EC, DeBeer BB, Silvia PJ, Gratz KL, Calhoun PS, Beckham JC and Morissette SB (2017) ADHD and nonsuicidal self-injury in male veterans with and without PTSD. *Psychiatry Research* 252, 161–163.
- Klonsky ED (2011) Non-suicidal self-injury in United States adults: prevalence, sociodemographics, topography and functions. *Psychological Medicine* 41, 1981–1986.
- Klonsky ED, Oltmanns TF and Turkheimer E (2003) Deliberate self-harm in a nonclinical population: prevalence and psychological correlates. *The American Journal of Psychiatry* 160, 1501–1508.
- Kuehn BM (2009) Soldier suicide rates continue to rise. *JAMA* 301, 1111–1113.
- Muehlenkamp JJ and Brausch AM (2016) Reconsidering criterion A for the diagnosis of non-suicidal self-injury disorder. *Journal of Psychopathology and Behavioral Assessment* 38, 547–558.
- Muehlenkamp JJ and Gutierrez PM (2007) Risk for suicide attempts among adolescents who engage in non-suicidal self-injury. *Archives of Suicide Research* 11, 69–82.
- Muehlenkamp JJ, Claes L, Havertape L and Plener PL (2012a) International prevalence of adolescent non-suicidal self-injury and deliberate self-harm. *Child & Adolescent Psychiatry & Mental Health* 6, 1–9.
- Muehlenkamp JJ, Peat CM, Claes L and Smits D (2012b) Self-injury and disordered eating: expressing emotion dysregulation through the body. *Suicide and Life-Threatening Behavior* 42, 416–425.
- Nock MK (2016) Recent and needed advances in the understanding, prediction, and prevention of suicidal behavior. *Depression and Anxiety* 33, 460–463.
- Nock MK, Joiner TE, Gordon KH, Lloyd-Richardson E and Prinstein MJ (2006) Non-suicidal self-injury among adolescents: diagnostic correlates and relation to suicide attempts. *Psychiatry Research* 144, 65–72.
- Nock MK, Hwang I, Sampson N, Kessler RC, Angermeyer M, Beautrais A, Borges G, Bromet E, Bruffaerts R, de Girolamo G, de Graaf R, Florescu S, Gureje O, Haro JM, Hu C, Huan Y, Karam EG, Kawakami N, Kovess V, Levinson D, Posada-Villa J, Sagar R, Tomov T, Viana MC and Williams DR (2009) Cross-national analysis of the associations among mental disorders and suicidal behavior: findings from the WHO World Mental Health Surveys. *PLoS Medicine* 6, e1000123.
- Nock MK, Hwang I, Sampson NA and Kessler RC (2010) Mental disorders, comorbidity and suicidal behavior: results from the National Comorbidity Survey Replication. *Molecular Psychiatry* 15, 868–876.
- Nock MK, Deming CA, Fullerton CS, Gilman SE, Goldenberg M, Kessler RC, McCarroll JE, McLaughlin KA, Peterson C, Schoenbaum M, Stanley B and Ursano RJ (2013a) Suicide among soldiers: a review of psychosocial risk and protective factors. *Psychiatry* 76, 97–125.
- Nock MK, Green JG, Hwang I, McLaughlin KA, Sampson NA, Zaslavsky AM and Kessler RC (2013b) Prevalence, correlates, and treatment of lifetime suicidal behavior among adolescents: results from the National Comorbidity Survey Replication Adolescent Supplement. *JAMA Psychiatry* 70, 300–310.
- Nock MK, Stein MB, Heeringa SG, Ursano RJ, Colpe LJ, Fullerton CS, Hwang I, Naifeh JA, Sampson NA, Schoenbaum M, Zaslavsky AM, Kessler RC and Army STARRS Collaborators (2014) Prevalence and correlates of suicidal behavior among soldiers: results from the Army Study to Assess Risk and Resilience in Servicemembers (Army STARRS). *JAMA Psychiatry* 71, 514–522.
- Nock MK, Millner AJ, Joiner TE, Gutierrez PM, Han G, Hwang I, King A, Naifeh JA, Sampson NA, Zaslavsky AM, Stein MB, Ursano RJ and Kessler RC (2018) Risk factors for the transition from suicide ideation to suicide attempt: results from the Army Study to Assess Risk and Resilience in Servicemembers (Army STARRS). *Journal of Abnormal Psychology* 127, 139–149.
- Plener PL, Allroggen M, Kapusta ND, Brähler E, Fegert JM and Groschwitz RC (2016) The prevalence of nonsuicidal self-injury (NSSI) in a representative sample of the German population. *BMC Psychiatry* 16, 1–7.
- Posner K, Brown GK, Stanley B, Brent DA, Yershova KV, Oquendo MA, Currier GW, Melvin GA, Greenhill L, Shen S and Mann JJ (2011) The Columbia-suicide severity rating scale: initial validity and internal consistency findings from three multisite studies with adolescents and adults. *American Journal of Psychiatry* 168, 1266–1277.
- Ribeiro JD, Franklin JC, Fox KR, Bentley KH, Kleiman EM, Chang BP and Nock MK (2016) Self-injurious thoughts and behaviors as risk factors for future suicide ideation, attempts, and death: a meta-analysis of longitudinal studies. *Psychological Medicine* 46, 225–236.
- Schoenbaum M, Kessler RC, Gilman SE, Colpe LJ, Heeringa SG, Stein MB, Ursano RJ, Cox KL and Army STARRS Collaborators (2014) Predictors of suicide and accident death in the Army Study to Assess Risk and Resilience in Servicemembers (Army STARRS): results from the Army Study to Assess Risk and Resilience in Servicemembers (Army STARRS). *JAMA Psychiatry* 71, 493–503.
- Shelf L, Fruchter E, Spiegel DO, Shoval G, Mann JJ and Zalsman G (2014) Characteristics of soldiers with self-harm in the Israeli Defense Forces. *Archives of Suicide Research* 18, 410–418.
- Singer JD and Willett JB (2003) *Applied Longitudinal Data Analysis: Modeling Change and Event Occurrence*, New York, NY, USA: Oxford University Press.

- Sobell LC and Sobell MB** (1992) Timeline follow-back: a technique for assessing self-reported alcohol consumption. In Litten RZ and Allen JP (eds), *Measuring Alcohol Consumption*. Totowa, NJ: Humana Press, pp. 41–72.
- Turner BJ, Yiu A, Layden BK, Claes L, Zaitsoff S and Chapman AL** (2015) Temporal associations between disordered eating and nonsuicidal self-injury: examining symptom overlap over 1 year. *Behavior Therapy* **46**, 125–138.
- Ursano RJ, Colpe LJ, Heeringa SG, Kessler RC, Schoenbaum M, Stein MB and Army STARRS collaborators** (2014) The Army Study to Assess Risk and Resilience in Servicemembers (Army STARRS). *Psychiatry* **77**, 107–119.
- Warner CH, Appenzeller GN, Parker JR, Warner CM and Hoge CW** (2011) Effectiveness of mental health screening and coordination of in-theater care prior to deployment to Iraq: a cohort study. *American Journal of Psychiatry* **168**, 378–385.
- Weathers FW, Litz BT, Keane TM, Palmieri PA, Marx BP and Schnurr PP** (2013) *The PTSD Checklist for DSM-5 (PCL-5)*. Washington, DC: National Center for PTSD.
- Whitlock J, Muehlenkamp JJ and Eckenrode J** (2008) Variation in nonsuicidal self-injury: identification and features of latent classes in a college population of emerging adults. *Journal of Clinical Child and Adolescent Psychology* **37**, 725–735.
- Whitlock J, Muehlenkamp J, Purington A, Eckenrode J, Barreira P, Baral Abrams G, Marchell T, Kress V, Girard K, Chin C and Knox K** (2011) Nonsuicidal self-injury in a college population: general trends and sex differences. *Journal of American College Health* **59**, 691–698.
- Whitlock JL and Selekmán M** (2014) Non-suicidal self-injury across the lifespan. In Nock M (ed.), *Oxford Handbook of Suicide and Self-Injury*. Oxford Library of Psychology: Oxford University Press.
- Wilkinson P, Kelvin R, Roberts C, Dubicka B and Goodyer I** (2011) Clinical and psychosocial predictors of suicide attempts and nonsuicidal self-injury in the Adolescent Depression Antidepressants and Psychotherapy Trial (ADAPT). *American Journal of Psychiatry* **168**, 495–501.
- Zetterqvist M, Lundh LG, Dahlström and Svedin CG** (2013) Prevalence and function of non-suicidal self-injury (NSSI) in a community sample of adolescents, using suggested DSM-5 criteria for a potential NSSI disorder. *Journal of Abnormal Child Psychology* **41**, 759–773.